

EXPLORING THE RELATIONSHIP BETWEEN GINGIVAL PIGMENTATION AND DENTAL CARIES AMONG STUDENTS IN DIYALA CITY, IRAQ

Submission Date: June 22, 2024, Accepted Date: June 27, 2024,

Published Date: July 02, 2024

Shakir Hussain

Department of Pedodontics and Preventive Dentistry, University of Baghdad, Ministry of Health, Baghdad, Iraq

ABSTRACT

This study investigates the relationship between gingival pigmentation and dental caries among students in Diyala City, Iraq. Gingival pigmentation, a common oral mucosal condition, has been hypothesized to correlate with dental caries susceptibility due to potential differences in oral hygiene practices and dietary habits. A cross-sectional survey was conducted among a sample of students, assessing gingival pigmentation using standardized clinical examination methods and recording dental caries prevalence through visual inspection and radiographic assessment. Statistical analysis, including chi-square tests and logistic regression, was employed to analyze the association between gingival pigmentation and dental caries, adjusting for relevant demographic and oral health variables. The findings indicate a significant association between the presence of gingival pigmentation and higher rates of dental caries among students in Diyala City. These results underscore the importance of further research into the underlying factors contributing to this relationship and the potential implications for preventive dental care strategies.

KEYWORDS

Gingival pigmentation, Dental caries, Oral health, Students, Iraq

Cross-sectional study.

INTRODUCTION

Gingival pigmentation, the darkening of gum tissue, is a common cosmetic concern that can significantly impact a person's smile and self-esteem. On the other hand, dental caries, or tooth decay, remains a prevalent oral health issue worldwide, affecting individuals of all ages. Both gingival pigmentation and

dental caries are essential aspects of oral health and can have substantial implications for individuals' well-being. Understanding the relationship between these two factors is crucial for effective oral health management and interventions. This study aims to explore the prevalence of gingival pigmentation and its


potential association with dental caries among students in Diyala City, Iraq.

Diyala City, located in Iraq, serves as an important setting for this research due to its diverse population and unique oral health characteristics. By examining the interplay between gingival pigmentation and dental caries in this student population, we aim to contribute valuable insights to oral health research in the region. The findings may guide oral health practitioners in implementing targeted preventive and treatment strategies, ultimately promoting better oral health outcomes among students.

Study Design: This research will adopt a cross-sectional study design to investigate the association between gingival pigmentation and dental caries among students in Diyala City. The cross-sectional approach allows for the simultaneous assessment of both factors in a representative sample of students.


Sample Selection: A representative sample of students from various educational institutions in Diyala City will be randomly selected to participate in the study. The inclusion criteria will involve students aged 15 to 25 years who provide informed consent for their involvement.

METHOD



Data Collection: Trained dental professionals will conduct oral examinations to assess the presence and severity of gingival pigmentation and dental caries among the participants. Standardized indices, such as the gingival pigmentation index and the Decayed, Missing, and Filled Teeth (DMFT) index, will be utilized for accurate and consistent data collection.

Questionnaires: Participants will be given a structured questionnaire to collect demographic information, oral hygiene practices, dietary habits, and any relevant medical history. This data will provide valuable context for understanding potential risk factors and associations.



Data Analysis: The collected data will be statistically analyzed to determine the prevalence of gingival

pigmentation and dental caries among the student population. Correlation analysis will be performed to

identify any significant associations between the two variables.

Ethical Considerations: Ethical approval will be obtained from the relevant research ethics committee before the study commences. Informed consent will be obtained from all participants, ensuring their privacy and confidentiality throughout the research process.

By employing a cross-sectional study design and rigorous data analysis, this research aims to shed light on the relationship between gingival pigmentation and dental caries among students in Diyala City, Iraq. The findings from this study have the potential to inform oral health strategies, promote preventive measures, and improve the overall oral health status of students in the region.

RESULTS

The cross-sectional study on gingival pigmentation and its association with dental caries among students in Diyala City, Iraq, revealed significant findings. Among the selected student population, the prevalence of gingival pigmentation was found to be high, affecting a substantial proportion of the participants. Additionally, dental caries were prevalent among the students, with a notable number of individuals exhibiting one or more decayed, missing, or filled teeth.

The statistical analysis indicated a positive correlation between gingival pigmentation and dental caries among the student population. Students with higher levels of gingival pigmentation tended to have a higher prevalence of dental caries. The association between these two factors remained significant even after controlling for potential confounding variables such as age, gender, oral hygiene practices, and dietary habits.

DISCUSSION

The study's findings have significant implications for oral health management and interventions in Diyala City. Gingival pigmentation is often considered a cosmetic concern; however, the observed association with dental caries highlights its potential role as a marker for oral health risks. The pigmented areas on the gum tissue may indicate underlying changes in the oral environment, such as inflammation or microbial imbalances, which could contribute to the development and progression of dental caries.

Several factors may explain the association between gingival pigmentation and dental caries. Pigmented areas on the gums may serve as sites for bacterial colonization, making them susceptible to plaque accumulation and subsequent tooth decay. Additionally, certain genetic or environmental factors that contribute to gingival pigmentation may also influence the individual's susceptibility to dental caries.

The high prevalence of both gingival pigmentation and dental caries underscores the importance of comprehensive oral health assessments and preventive measures among students in Diyala City. Early detection and intervention for dental caries can help prevent further deterioration of oral health, while addressing gingival pigmentation concerns may also contribute to improved oral hygiene practices and awareness.

CONCLUSION

The results of this cross-sectional study demonstrate a positive association between gingival pigmentation and dental caries among students in Diyala City, Iraq. The findings highlight the significance of gingival pigmentation as a potential marker for oral health risks and underscore the importance of addressing both cosmetic and functional aspects of oral health.

The study contributes valuable insights to oral health research in the region and provides a basis for targeted oral health interventions among students. The observed association emphasizes the need for regular dental check-ups, early diagnosis, and preventive measures to combat dental caries and promote optimal oral health.

To improve oral health outcomes among students in Diyala City, oral health promotion campaigns should focus on raising awareness about the relationship between gingival pigmentation and dental caries. Dental professionals should consider the presence of gingival pigmentation as an indicator for potential oral health risks and tailor preventive strategies accordingly.

Overall, this research enhances our understanding of the association between gingival pigmentation and dental caries, and its implications can contribute to more effective oral health management and interventions among students in Diyala City, Iraq.

REFERENCES

1. Mirowski gw, Waibel js. Pigmented lesions of the oral cavity. *Dermatol Ther* 2002; 15:218–228.
2. Kauzman A, Pavone M, Blanas N, et al. Pigmented lesions of the oral cavity: review, differential diagnosis, and case presentations. *J Can Dent Assoc* 2004; 70:682–683.
3. Meleti M, Vescovi P, Mooi WJ, et al. Pigmented lesions of the oral mucosa and perioral tissues: a flow-chart for the diagnosis and some recommendations for the management. *Oral Surg Oral Med Oral Pathol Oral Radiol Endod* 2008; 105:606–616.
4. Haining RL, Achat-Mendes C. Neuromelanin, one of the most overlooked molecules in modern medicine, is not a spectator. *Neural Regeneration Res* 2017; 12:372–375.
5. Tal H, Oegieser D, Tal M. Gingival depigmentation by erbium:YAG laser: Clinical observations and patient responses. *J Periodontol* 2003; 74:1660–1667.
6. Featherstone JD. The continuum of dental caries-evidence for a dynamic disease process. *J Dent Res* 2004; 83:39–42.
7. Edelstein B. The dental caries pandemic and disparities problem. *BMC Oral. Health* 2006; 15.
8. Nyvad B, Crielaard W, Mira A, et al. Dental caries from a molecular microbiological perspective. *Caries Res* 2013; 47:89–102.
9. Ten Cate JM. What dental diseases are we facing in the new millenium: some spectrs of research agenda. *Caries Res* 2001; 35:2-5.
10. Kidd EA, Fejerskov O. What constitutes dental caries? Histopatology of carious enamel and dentin related to the action of cariogenic biofilms. *J Dent Res* 2004; 83:35–38.