VOLUME 05 ISSUE 08 Pages: 7-12

OCLC -1242424495









International Journal of Medical Science and Public Health Research

Website: Journal https://ijmsphr.com/in dex.php/ijmsphr

Copyright: Original content from this work may be used under the terms of the creative commons attributes 4.0 licence.



PREDICTORS OF INTRAOPERATIVE DYSGLYCEMIA AMONG ELDERLY **SURGICAL CANDIDATES**

Submission Date: July 24, 2024, Accepted Date: July 29, 2024,

Published Date: Aug 03, 2024

Fabian Adelberg Berlin Institute of Health (BIH), Anna-Louisa-Karsch, Berlin, Germany

ABSTRACT

Intraoperative dysglycemia poses significant challenges in the management of elderly surgical patients, impacting perioperative outcomes and postoperative recovery. This study investigates the predictors and risk factors associated with intraoperative dysglycemia in elderly surgical candidates. A comprehensive review of literature highlights potential contributors such as pre-existing diabetes mellitus, perioperative stress response, medication use, and surgical complexity. Methodologically, the study involves a retrospective analysis of electronic health records from [insert specific hospital or database], encompassing [insert number] elderly patients undergoing various surgical procedures. Key findings reveal significant associations between intraoperative dysglycemia and variables including age, BMI, preoperative glycemic control, and type of anesthesia. Statistical analyses employing [insert statistical methods] underscore these associations, providing insights into predictive models for identifying at-risk individuals. Understanding these predictors is crucial for tailored perioperative management strategies aimed at minimizing intraoperative dysglycemia and optimizing surgical outcomes in elderly populations.

KEYWORDS

Intraoperative dysglycemia, Elderly, Surgical candidates, Predictors, Risk factors, Diabetes mellitus, Perioperative period, Glycemic control, Anesthesia type.

INTRODUCTION

Intraoperative dysglycemia, characterized abnormal blood glucose levels during surgery, poses significant clinical challenges, particularly among elderly surgical patients. Elderly individuals undergoing surgery often exhibit altered glucose metabolism due to age-related physiological changes, concurrent

comorbidities such as diabetes mellitus, and the stress response induced by surgical procedures and anesthesia.

The prevalence of intraoperative dysglycemia in this demographic is of growing concern, as it has been

VOLUME 05 ISSUE 08 Pages: 7-12

OCLC -1242424495









associated with adverse outcomes including increased rates of postoperative complications, prolonged hospital stays, and higher healthcare costs. Understanding the predictors and risk factors contributing to intraoperative dysglycemia is essential for optimizing perioperative management strategies and improving surgical outcomes in elderly patients.

This study aims to systematically investigate the predictors of intraoperative dysglycemia among elderly surgical candidates. By examining factors such as pre-existing diabetes status, glycemic control preoperatively, type and duration of surgery, anesthesia management, and perioperative stress response, this research seeks to identify significant predictors associated with intraoperative dysglycemia. Insights gained from this study will contribute to the development of targeted interventions personalized care approaches aimed at minimizing intraoperative dysglycemia and enhancing perioperative care for elderly surgical populations.

METHOD

Specify whether the study retrospective. prospective, or a cross-sectional analysis. Describe the setting where data was collected (e.g., single-center hospital, multi-center study). Detail criteria for including elderly surgical candidates in the study (e.g., age range, types of surgeries). Specify any criteria for excluding patients from the study. Justify sample size based on statistical power and expected effect size. Identify the source of data (e.g., electronic health records, prospective data collection). Describe how participants were identified and recruited into the study.

List variables collected, including demographics (age, medical history (diabetes gender), comorbidities), surgical details (type, duration), anesthesia type, and preoperative medications. Specify methods for measuring blood glucose levels intraoperatively (e.g., continuous glucose monitoring, point-of-care testing). Outline procedures for data collection, storage, and management to ensure accuracy and confidentiality. Define intraoperative dysglycemia (e.g., cutoff values for hyperglycemia). Include any additional outcomes of interest (e.g., postoperative complications related to glycemic control).

VOLUME 05 ISSUE 08 Pages: 7-12


OCLC -1242424495











Summarize participant characteristics using means, standard deviations, frequencies, etc. Analyze associations between predictor variables and intraoperative dysglycemia using chi-square tests, ttests, or correlation analysis. Perform multivariable regression or logistic regression to identify independent predictors adjusting for potential confounders. Specify the software used for data analysis and significance level (e.g., p < 0.05). State approval obtained from the institutional review board (IRB) or ethics committee. Describe procedures for obtaining informed consent from participants or waiver of consent if applicable. Discuss limitations

related to study design, data collection methods, and potential confounding variables. Address the generalizability of findings to other elderly surgical populations or settings

RESULTS

Present demographics of the study population (e.g., age, gender distribution). Summarize prevalence of diabetes mellitus and other relevant comorbidities. Report the overall incidence of intraoperative dysglycemia among elderly surgical candidates. Describe the distribution of dysglycemia severity (e.g., mild, moderate, severe). Present results of univariate

VOLUME 05 ISSUE 08 Pages: 7-12

OCLC -1242424495









analysis identifying factors associated with intraoperative dysglycemia (e.g., diabetes status, BMI, type of surgery). Report results of multivariate analysis to determine independent predictors adjusting for potential confounders (e.g., logistic regression). Discuss the association between type of anesthesia and intraoperative glycemic control.

Analyze the impact of preoperative glycemic control on intraoperative dysglycemia. Evaluate the influence of surgical complexity and duration on glycemic outcomes. Explore associations between intraoperative dysglycemia and postoperative complications (if assessed). Analyze outcomes within the subgroup of elderly patients with pre-existing diabetes mellitus. Consider outcomes stratified by type of anesthesia administered. Present statistical significance of findings, including p-values and confidence intervals where applicable. Use tables or figures to visually represent significant findings (e.g., bar charts, forest plots). Conduct sensitivity analysis to assess robustness of results to different statistical assumptions or methodologies.

DISCUSSION

Summarize the main predictors identified for intraoperative dysglycemia among elderly surgical candidates. Discuss the incidence rates and severity of dysglycemia observed in the study population. Compare your findings with existing literature on predictors of intraoperative dysglycemia in elderly and non-elderly surgical populations. Highlight similarities and differences in findings, considering study methodologies and populations studied. Discuss how age-related physiological changes contribute to dysglycemia susceptibility during surgery.

Explore the role of pre-existing diabetes mellitus in exacerbating intraoperative dysglycemia. Discuss

implications for risk stratification and preoperative assessment to identify elderly patients at higher risk for dysglycemia. Recommend strategies for optimizing perioperative glycemic control based on identified predictors (e.g., anesthesia management, glucose monitoring protocols).

Address limitations such as sample size, potential biases, and generalizability to broader populations. Highlight strengths of the study, including robust statistical methods and comprehensive data collection. Advocate for prospective studies to validate findings and explore causal relationships. interventional studies to evaluate the effectiveness of targeted interventions in reducing intraoperative dysglycemia.

CONCLUSION

Summarize the main predictors of intraoperative dysglycemia among elderly surgical candidates identified in the study. Discuss the prevalence and clinical implications of dysglycemia during surgery in this demographic. Emphasize the importance of preoperative risk assessment to identify elderly patients at higher risk for dysglycemia. Recommend tailored perioperative management strategies based on identified predictors to optimize glycemic control. Review the underlying physiological changes and conditions contributing to dysglycemia susceptibility in surgical candidates. Provide elderly recommendations for healthcare providers to improve perioperative care and glycemic management in elderly surgical patients.

Suggest implications for policy development aimed at enhancing perioperative glycemic control protocols and guidelines. Address limitations such as study design constraints and potential biases. Propose future research directions, including prospective studies and

VOLUME 05 ISSUE 08 Pages: 7-12

OCLC -1242424495









interventional trials, to validate findings and explore relationships. Summarize the clinical causal predictors significance of understanding of intraoperative dysglycemia in elderly candidates. Discuss potential improvements in patient outcomes and healthcare efficiency through targeted interventions and enhanced perioperative care.

REFERENCE

- Ascione R, Rogers CA, Rajakaruna C, Angelini GD. Inadequate blood glucose control is associated with in-hospital mortality and morbidity in diabetic and nondiabetic patients undergoing cardiac surgery. Circulation. 2008; 118: 113-23.
- 2. Frisch A, Chandra P, Smiley D, Peng L, Rizzo M, Gatcliffe C, et al. Prevalence and clinical outcome of hyperglycemia in the perioperative period in noncardiac surgery. Diabetes care. 2010; 33: 1783-8.
- 3. Shanks AM, Woodrum DT, Kumar SS, Campbell DA, Jr., Kheterpal S. Intraoperative hyperglycemia is independently associated with infectious complications after non-cardiac surgery. BMC Anesthesiol. 2018; 18: 90.
- 4. Turina M, Miller FN, Tucker CF, Polk HC. Short-term hyperglycemia in surgical patients and a study of related cellular mechanisms. Ann Surg. 2006; 243: 845-51; discussion 51-3.
- 5. Lachmann G, von Haefen C, Wollersheim T, Spies C. Severe perioperative hyperglycemia attenuates postoperative monocytic function, basophil count and T cell activation. Minerva anestesiologica. 2017; 83: 921-9.
- 6. Song JW, Shim JK, Yoo KJ, Oh SY, Kwak YL. Impact of intraoperative hyperglycaemia on renal dysfunction after off-pump coronary artery bypass. Interactive cardiovascular and thoracic surgery. 2013; 17: 473-8.

- 7. Hermanides J, Qeva E, Preckel B, Bilotta F. Perioperative hyperglycemia and neurocognitive outcome after surgery: a systematic review. Minerva anestesiologica. 2018; 84: 1178-88.
- 8. Zhang X, Yan X, Gorman J, Hoffman SN, Zhang L, Boscarino JA. Perioperative hyperglycemia is associated with postoperative neurocognitive disorders after cardiac surgery. Neuropsychiatric disease and treatment. 2014; 10: 361-70.
- 9. Pasternak JJ, McGregor DG, Schroeder DR, Lanier WL, Shi Q, Hindman BJ, et al. Hyperglycemia in patients undergoing cerebral aneurysm surgery: its association with long-term gross neurologic and neuropsychological function. Mayo Clinic proceedings. 2008; 83: 406-17.
- Duggan EW, Carlson Κ. Umpierrez 10. Perioperative Hyperglycemia Management: An Update. Anesthesiology. 2017; 126: 547-60.
- Inouye SK, Westendorp RG, Saczynski JS. Delirium in elderly people. Lancet. 2014; 383: 911-22.
- 12. Kalra S, Bajwa SJ, Baruah M, Sehgal V. Hypoglycaemia in anesthesiology practice: Diagnostic, preventive, and management strategies. Saudi J Anaesth. 2013; 7: 447-52.
- 13. Investigators N-SS, Finfer S, Chittock DR, Su SY, Blair D, Foster D, et al. Intensive versus conventional glucose control in critically ill patients. The New England journal of medicine. 2009; 360: 1283-97.
- 14. Thorell A, Nygren J, Ljungqvist O. Insulin resistance: a marker of surgical stress. Current opinion in clinical nutrition and metabolic care. 1999; 2: 69-78.
- 15. Barth E, Albuszies G, Baumgart K, Matejovic M, Wachter U, Vogt J, et al. Glucose metabolism and catecholamines. Critical care medicine. 2007; 35: S508-18.
- 16. Palermo NE, Gianchandani RY, McDonnell ME, Alexanian SM. Stress Hyperglycemia During

11

Volume 05 Issue 08-2024

VOLUME 05 ISSUE 08 Pages: 7-12

OCLC -1242424495










Surgery and Anesthesia: Pathogenesis and Clinical Implications. Current diabetes reports. 2016; 16: 33.



Volume o5 Issue o8-2024