



OPEN ACCESS

SUBMITTED 15 June 2025

ACCEPTED 11 July 2025

PUBLISHED 13 August 2025

VOLUME Vol.06 Issue08 2025

CITATION

F.A. Kandova, & N.N.Khabibova. (2025). Impact of hormonal fluctuations on oral conditions in reproductive-age women. International Journal of Medical Science and Public Health Research, 6(08), 6–10.

<https://doi.org/10.37547/ijmsphr/Volume06Issue08-02>

COPYRIGHT

© 2025 Original content from this work may be used under the terms of the creative commons attributes 4.0 License.

Impact of hormonal fluctuations on oral conditions in reproductive-age women

F.A. Kandova

Bukhara Innovative University of Education and Medicine, Bukhara, Uzbekistan

N.N.Khabibova

Bukhara State Medical Institute named after Abu Ali ibn Sino, Bukhara, Uzbekistan

Abstract: Hormonal fluctuations throughout the reproductive years play a crucial role in women's systemic and oral health. Estrogen and progesterone variations during menstrual cycles, contraceptive use, and preconceptional phases can significantly influence the condition of the oral mucosa, gingival tissues, and salivary composition. These hormonal changes may lead to clinical manifestations such as gingival inflammation, bleeding, swelling, and mucosal discomfort, even in the absence of significant plaque accumulation. This article reviews the physiological mechanisms linking hormonal activity to oral conditions, outlines common hormone-related oral pathologies, and presents current approaches for screening and managing such issues in women preparing for pregnancy. Emphasis is placed on the importance of early diagnosis and interdisciplinary care in the context of reproductive planning.

Keywords: Hormonal fluctuations, oral health, reproductive-age women, gingival inflammation, preconception care, estrogen, progesterone.

Introduction: Women's health during the reproductive years is deeply influenced by cyclical hormonal fluctuations that impact various physiological systems, including the oral cavity. The two primary female sex hormones—estrogen and progesterone—are known to modulate immune response, vascular permeability, and connective tissue metabolism. These effects become particularly relevant during specific phases such as the menstrual cycle, hormonal contraceptive use, and the preconception period. While much focus is placed on

the reproductive and emotional aspects of hormonal variation, its implications for oral health remain underrecognized.

Emerging evidence suggests that hormonal changes can increase the risk of inflammatory conditions in the oral cavity, particularly in the gingival tissues. Women of reproductive age commonly report symptoms such as gum tenderness, bleeding, and swelling—symptoms that are not always attributable to poor oral hygiene alone but rather reflect the underlying endocrine environment. Furthermore, hormonal shifts can influence the composition of saliva, alter oral microbiota, and exacerbate preexisting periodontal conditions.

In the context of pregnancy planning, the oral cavity requires special attention. Poor oral health prior to conception has been linked to adverse outcomes such as preterm birth, preeclampsia, and low birth weight. As such, identifying and managing hormonally influenced oral conditions during the preconception phase is crucial for safeguarding both maternal and fetal health.

Purpose of the research

This article aims to explore the relationship between hormonal fluctuations and oral conditions in women of reproductive age. It focuses on the biological mechanisms involved, describes the most common hormone-associated oral manifestations, and outlines strategies for effective screening and preventive care tailored to women preparing for pregnancy.

Hormonal Fluctuations and Their Impact on Oral Tissues

Hormonal fluctuations are a hallmark of a woman's reproductive life, manifesting during the menstrual cycle, pregnancy, lactation, and in response to hormonal contraceptives. These hormonal variations, particularly in estrogen and progesterone levels, have profound effects on the oral mucosa, gingival vasculature, and immune regulation. As a result, even in women with good oral hygiene, hormone-related changes may cause exaggerated tissue responses to minor irritants such as dental plaque.

Estrogen promotes increased vascularization and epithelial proliferation in oral tissues, leading to enhanced capillary permeability and edema. These changes contribute to gingival swelling, increased bleeding on probing, and greater susceptibility to irritation. On the other hand, progesterone reduces

collagen production and affects neutrophil chemotaxis, impairing the host's ability to respond effectively to microbial invasion [1].

Several phases of hormonal activity are particularly relevant:

Menstrual cycle: Many women experience gingival discomfort or transient inflammation during the luteal phase, when progesterone levels peak.

Hormonal contraceptive use: Long-term exposure to synthetic estrogens and progestins can mimic pregnancy-like changes in gingival tissues, often leading to gingival hyperplasia or exacerbation of chronic periodontitis.

Preconception period: Hormonal priming of the endometrium and associated systemic changes may cause subclinical shifts in oral immunity, paving the way for latent gingival inflammation or mucosal dryness.

Moreover, these hormonal effects can disturb the salivary gland function, reducing flow rate and altering its buffering capacity and antimicrobial properties. Such changes increase vulnerability to caries, oral candidiasis, and halitosis [2].

Understanding the interaction between systemic hormones and the oral cavity is essential for early identification and personalized management of oral conditions in reproductive-age women. Dental practitioners should be aware of these patterns and incorporate hormonal history into their diagnostic evaluations, especially in women planning pregnancy.

Relationship Between Hormonal Changes and Oral Diseases

A growing body of clinical and epidemiological evidence highlights the close association between hormonal fluctuations and the prevalence of various oral diseases in women of reproductive age. These conditions are often classified as hormone-responsive oral disorders, as they tend to emerge or worsen in response to cyclical endocrine changes.

One of the most common manifestations is gingivitis, particularly menstrual cycle-associated gingivitis and pregnancy gingivitis. Studies show that elevated progesterone levels increase gingival vascular permeability and prostaglandin production, which heightens inflammatory responses to plaque. As a result, women may experience reddening, swelling, and spontaneous bleeding of the gums, often without

proportional plaque accumulation [3].

Gingival overgrowth (also referred to as gingival hyperplasia) is another condition associated with hormonal influence, especially among women using oral contraceptives or undergoing hormonal therapy. This condition not only affects aesthetics but can also impair proper oral hygiene, leading to further periodontal complications [4].

Periodontitis, a chronic inflammatory condition affecting the supporting structures of the teeth, may be aggravated by hormonal changes due to immune modulation. Hormonal imbalance can enhance the activity of cytokines such as IL-1 β , IL-6, and TNF- α , which contribute to tissue destruction and alveolar bone loss. When left untreated in the preconception period, periodontitis may increase the risk of adverse

pregnancy outcomes such as preterm birth and low birth weight [5].

Oral candidiasis also tends to be more frequent in hormonally unstable periods. Progesterone-induced immunosuppression and altered salivary composition may create favorable conditions for *Candida albicans* colonization.

Burning mouth syndrome, metallic taste sensations, and xerostomia (dry mouth) are additional conditions that are reported more frequently in women experiencing hormonal imbalances, especially in those with underlying thyroid or endocrine disorders [6].

The table below summarizes the most common hormone-associated oral diseases and their typical clinical features:

Table 1. Common hormone-related oral conditions in reproductive-age women

Condition	Hormonal Trigger	Clinical Manifestations
Gingivitis	Menstrual cycle, pregnancy, OCs	Swelling, redness, bleeding
Gingival hyperplasia	Oral contraceptive use	Tissue enlargement, poor plaque control
Periodontitis	Chronic hormonal imbalance	Bone loss, pocket formation, tooth mobility
Oral candidiasis	Immunosuppression, altered saliva	White plaques, burning, discomfort
Xerostomia	Estrogen/progesterone variation	Dry mouth, sticky feeling, speech difficulty

Diagnostic Methods for Assessing Hormone-Linked Oral Changes

Timely and accurate diagnosis of hormone-related oral conditions in women preparing for pregnancy is essential to prevent complications during gestation and ensure optimal maternal health. These conditions may not always be evident through conventional clinical examination alone, and therefore a combination of targeted diagnostic tools is recommended.

Comprehensive clinical examination

Initial diagnosis begins with thorough inspection of the oral cavity. Dentists should assess gingival color, texture, bleeding on probing, tissue enlargement, and

any mucosal lesions. A detailed menstrual and contraceptive history should also be recorded to correlate symptoms with hormonal phases [1].

Periodontal indices

Validated indices such as the Gingival Index (GI), Plaque Index (PI), Bleeding on Probing (BOP), and Periodontal Probing Depth (PPD) are essential tools to quantify inflammation, plaque accumulation, and periodontal pocketing. Tracking these values over time can help detect hormonal influence on disease progression [2].

Levels of secretory immunoglobulin A (SIgA), lysozyme, IL-6, and TNF- α can signal mucosal immune activity.

Advanced salivary tests using enzyme-linked

immunosorbent assay (ELISA) or rapid strip tests are available for chairside screening [3].

Microbiological testing

Hormonal shifts can lead to changes in the oral microbiome. Collection of subgingival plaque and mucosal swabs for microbial culture or PCR (polymerase chain reaction) testing can help identify overgrowth of periodontopathogenic bacteria or fungal elements such as *Candida albicans* [4].

Imaging and visualization tools

In selected cases, intraoral cameras, fluorescence-based diagnostics (VELscope), or laser Doppler imaging may be employed to detect subtle mucosal or vascular changes that are associated with hormone-influenced tissue response [5].

Interdisciplinary collaboration

Dentists should consider referring patients to gynecologists or endocrinologists for hormonal profiling (estrogen, progesterone, prolactin, thyroid function tests) when oral signs persist despite conventional therapy.

Early detection of hormone-related oral disturbances allows for preventive intervention prior to pregnancy. Personalized treatment plans based on diagnostic findings can significantly reduce discomfort and improve overall reproductive outcomes.

CONCLUSION

Hormonal fluctuations in reproductive-age women have a profound impact on oral health, influencing tissue metabolism, immune responses, and microbial balance within the oral cavity. These endocrine shifts, while physiological, can predispose women to various oral conditions such as gingivitis, periodontitis, candidiasis, and xerostomia—even in the absence of poor hygiene or other systemic disease.

Recognizing the link between hormonal activity and oral manifestations is particularly crucial during the preconception period, as undiagnosed or untreated oral diseases may contribute to adverse pregnancy outcomes. A proactive approach involving early screening, personalized oral care, and interdisciplinary coordination between dental and medical professionals can ensure optimal maternal oral health and reduce the risk of pregnancy-related complications.

Incorporating hormonal considerations into routine dental assessments is not only clinically relevant but essential in the comprehensive care of women preparing for pregnancy.

REFERENCES

- Mariotti, A. (1994). Sex steroid hormones and cell dynamics in the periodontium. *Critical Reviews in Oral Biology & Medicine*, 5(1), 27–53. <https://doi.org/10.1177/10454411940050010201>
- Gursoy, M., Pajukanta, R., Sorsa, T., & Kononen, E. (2008). Clinical changes in periodontium during pregnancy and post-partum. *Journal of Clinical Periodontology*, 35(7), 576–583. <https://doi.org/10.1111/j.1600-051X.2008.01234.x>
- Kornman, K. S., & Loesche, W. J. (1982). The subgingival microbial flora during pregnancy. *Journal of Periodontal Research*, 17(2), 111–122. <https://doi.org/10.1111/j.1600-0765.1982.tb00092.x>
- Mealey, B. L., & Moritz, A. J. (2003). Hormonal influences: Effects of diabetes mellitus and endocrine dysfunction on the periodontium. *Periodontology 2000*, 32(1), 59–81. <https://doi.org/10.1046/j.0906-6713.2002.03206.x>
- Figuro, E., Carrillo-de-Albornoz, A., Martín, C., & Herrera, D. (2013). Effect of pregnancy on gingival inflammation in systemically healthy women: A systematic review. *Journal of Clinical Periodontology*, 40(5), 457–473. <https://doi.org/10.1111/jcpe.12071>
- Gürsoy, M., Könönen, E., Pradhan-Palikhe, P., et al. (2010). Salivary cytokines and the association of pregnancy with gingivitis. *Journal of Clinical Periodontology*, 37(6), 513–518. <https://doi.org/10.1111/j.1600-051X.2010.01561.x>
- Han, Y. W. (2011). Oral health and adverse pregnancy outcomes—What's next? *Journal of Dental Research*, 90(3), 289–293. <https://doi.org/10.1177/0022034510381905>
- Laine, M. A. (2002). Effect of pregnancy on periodontal and dental health. *Acta Odontologica Scandinavica*, 60(5), 257–264. <https://doi.org/10.1080/00016350260248210>
- Soell, M., Elkaim, R., Tenenbaum, H., & Davideau, J. L. (2002). Cathepsin L expression in healthy and diseased human gingiva: Effect of hormonal status. *Journal of Periodontology*, 73(4), 424–430. <https://doi.org/10.1902/jop.2002.73.4.424>

American Dental Association. (2021). Women's oral health: Hormonal influences and clinical care. Retrieved from <https://www.ada.org/resources/research/science-and-research-institute>